
Examining the usefulness of trophic level-based indicators in assessing ecosystem-level changes in the ocean

Lynne Shannon^{*1}, Marta Coll^{2,3}, Yunne Shin^{4,5}, Jodie Reed^{1,4}, Alida Bundy⁶, and Kristin Kleisner⁷

¹Marine Research Institute, University of Cape Town (Ma-Re, UCT) – Private Bag X3 Rondebosch 7701, South Africa

²Institut de Recherche pour le Développement (UMR EME 212) – Institut de recherche pour le développement [IRD] : UMR212 – France

³Institute of Marine Science (ICM-CSIC) – Spain

⁴Institut de Recherche pour le Développement (CRH, IRD) – CRH – Institut de Recherche pour le Développement, CRH, Research Unit MARBEC (UMR 248), avenue Jean Monnet, CS 30171,, France

⁵Université de Montpellier – – Université de Montpellier, Place Eugène Bataillon, Bâtiment 24, CC 093, 34095 Montpellier cedex 05, France, France

⁶Fisheries and Oceans Canada – Fisheries and Oceans Canada, Bedford Institute of Oceanography, 1 Challenger Drive, Dartmouth, NS B2Y 4A2, Canada, Canada

⁷Environmental Defense Fund, 18 Tremont St., Ste. 850, Boston, MA 02108 – Environmental Defense Fund, 18 Tremont St., Ste. 850, Boston, MA 02108, United States

Abstract

Trophic level (TL)-based indicators have been widely used to examine fishing impacts in aquatic ecosystems and induced biodiversity changes. However, there has been much debate concerning the use of landings data from commercial fisheries to calculate TL indicators because commercially caught species and their relative catches likely do not reflect the whole ecosystem and strongly depend on fishing strategies. Subsequent studies have started to look at survey-based and model-based TL indicators. Further, the use of a fixed TL per species in the calculation of TL-based indicators has been questioned, given that species' TLs vary with ontogeny, as well as over time and space. In this contribution, we report on two studies that contribute to this discussion: (i) a data-based extensive evaluation of a variety of trophic level indicators across nine well-studied marine ecosystems using model-based, survey-based, and catch-based TL indicators as well as detailed regional information, data on fishing history, fishing intensity, and environmental conditions, to examine how well TL indicators capture fishing effects at the community level of marine ecosystems; (ii) a fishing scenarios study using four different ecosystem modelling approaches in contrasted ecosystem case studies to examine the performance of TL-based indicators in tracking the effects of fishing pressure when using fixed species TL versus species' TL varying depending on the fishing scenarios. Results of the first study highlight that the differences observed between TL indicators' trends are dependent on the data source, and are not attributable to an intrinsic problem with TL-based indicators. All three data sources provide useful information about the structural

*Speaker

changes in the ecosystem as a result of fishing, but only model-based indicators were found to represent fishing impacts at the whole ecosystem level. From the second study, we conclude that overall, TL-based indicators calculated with variable species' TL have a greater capacity for detecting the effects of fishing pressure than when using fixed species' TL. Across TL-based indicators, survey-based TL indicators displayed the greatest consistency regarding the use of fixed or variable species' TL and the highest capacity for detecting fishing effects. This work supports the use of community-based indicators over catch-based indicators to explore the impacts of fishing on the structure of marine ecosystems.

Keywords: trophic level, fishing effects, indicators, fishing scenarios, trophic model